

Date: October 28, 2014

From: B.M. Wojek / Modified by A. Suter

E-Mail: andreas.suter@psi.ch

MUSRFIT plug-in for the calculation of the temperature dependence of $1/\lambda^2$ for various gap symmetries

This memo is intended to give a short summary of the background on which the GAPINTEGRALS plug-in for MUSRFIT [1] has been developed. The aim of this implementation is the efficient calculation of integrals of the form

$$I(T) = 1 + \frac{1}{\pi} \int_0^{2\pi} \int_{\Delta(\varphi, T)}^{\infty} \left(\frac{\partial f}{\partial E} \right) \frac{E}{\sqrt{E^2 - \Delta^2(\varphi, T)}} dE d\varphi, \quad (1)$$

where $f = (1 + \exp(E/k_B T))^{-1}$, like they appear e.g. in the theoretical temperature dependence of $1/\lambda^2$ [4]. In order not to do too many unnecessary function calls during the final numerical evaluation we simplify the integral (1) as far as possible analytically. The derivative of f is given by

$$\frac{\partial f}{\partial E} = -\frac{1}{k_B T} \frac{\exp(E/k_B T)}{(1 + \exp(E/k_B T))^2} = -\frac{1}{4k_B T} \frac{1}{\cosh^2(E/2k_B T)}. \quad (2)$$

Using (2) and doing the substitution $E'^2 = E^2 - \Delta^2(\varphi, T)$, equation (1) can be written as

$$\begin{aligned} I(T) &= 1 - \frac{1}{4\pi k_B T} \int_0^{2\pi} \int_{\Delta(\varphi, T)}^{\infty} \frac{1}{\cosh^2(E/2k_B T)} \frac{E}{\sqrt{E^2 - \Delta^2(\varphi, T)}} dE d\varphi \\ &= 1 - \frac{1}{4\pi k_B T} \int_0^{2\pi} \int_0^{\infty} \frac{1}{\cosh^2(\sqrt{E'^2 + \Delta^2(\varphi, T)}/2k_B T)} dE' d\varphi. \end{aligned} \quad (3)$$

Since a numerical integration should be performed and the function to be integrated is exponentially approaching zero for $E' \rightarrow \infty$ the infinite E' integration limit can be replaced by a cutoff energy E_c which has to be chosen big enough:

$$I(T) \simeq \tilde{I}(T) \equiv 1 - \frac{1}{4\pi k_B T} \int_0^{2\pi} \int_0^{E_c} \frac{1}{\cosh^2(\sqrt{E'^2 + \Delta^2(\varphi, T)}/2k_B T)} dE' d\varphi. \quad (4)$$

In the case that $\Delta^2(\varphi, T)$ is periodic in φ with a period of $\pi/2$ (valid for all gap symmetries implemented at the moment), it is enough to limit the φ -integration to one period and to multiply the result by 4:

$$\tilde{I}(T) = 1 - \frac{1}{\pi k_B T} \int_0^{\pi/2} \int_0^{E_c} \frac{1}{\cosh^2(\sqrt{E'^2 + \Delta^2(\varphi, T)}/2k_B T)} dE' d\varphi. \quad (5)$$

For the numerical integration we use algorithms of the CUBA library [2] which require to have a Riemann integral over the unit square. Therefore, we have to scale the integrand by the upper limits of the integrations. Note that E_c and $\pi/2$ (or in general the upper limit of the φ integration) are now treated as dimensionless scaling factors.

$$\tilde{I}(T) = 1 - \frac{E_c}{2k_B T} \int_0^{1\varphi} \int_0^{1E} \frac{1}{\cosh^2(\sqrt{(E_c E)^2 + \Delta^2(\frac{\pi}{2}\varphi, T)}/2k_B T)} dE d\varphi \quad (6)$$

Implemented gap functions and function calls from MUSRFIT

Currently the calculation of $\tilde{I}(T)$ is implemented for various gap functions. The temperature dependence of the gap functions is either given by Eq.(7) [3], or by Eq.(8) [4].

$$\Delta(\varphi, T) \simeq \Delta(\varphi, 0) \tanh \left[\frac{\pi k_B T_c}{\Delta_0} \sqrt{a_G \left(\frac{T_c}{T} - 1 \right)} \right] \quad (7)$$

with Δ_0 as given below, and a_G depends on the pairing state:

s-wave: $a_G = 1$ with $\Delta_0 = 1.76 k_B T_c$

d-wave: $a_G = 4/3$ with $\Delta_0 = 2.14 k_B T_c$

$$\Delta(\varphi, T) \simeq \Delta(\varphi) \tanh \left(1.82 \left(1.018 \left(\frac{T_c}{T} - 1 \right) \right)^{0.51} \right). \quad (8)$$

The GAPINTEGRALS plug-in calculates $\tilde{I}(T)$ for the following $\Delta(\varphi)$:

s-wave gap:

$$\Delta(\varphi) = \Delta_0 \quad (9)$$

MUSRFIT theory line: `userFcn libGapIntegrals TGapSWave 1 2 [3]`

(Parameters: T_c (K), Δ_0 (meV), [a_G (1)]. If a_G is given, the temperature dependence according to Eq.(7) will be used, otherwise Eq.(8) will be utilized.)

d-wave gap [5]:

$$\Delta(\varphi) = \Delta_0 \cos(2\varphi) \quad (10)$$

MUSRFIT theory line: `userFcn libGapIntegrals TGapDWave 1 2 [3]`

(Parameters: T_c (K), Δ_0 (meV), [a_G (1)]. If a_G is given, the temperature dependence according to Eq.(7) will be used, otherwise Eq.(8) will be utilized.)

non-monotonic d-wave gap [6]:

$$\Delta(\varphi) = \Delta_0 [a \cos(2\varphi) + (1 - a) \cos(6\varphi)] \quad (11)$$

MUSRFIT theory line: `userFcn libGapIntegrals TGapNonMonDWave1 1 2 3 [4]`

(Parameters: T_c (K), Δ_0 (meV), a (1), [a_G (1)]. If a_G is given, the temperature dependence according to Eq.(7) will be used, otherwise Eq.(8) will be utilized.)

non-monotonic d-wave gap [7]:

$$\Delta(\varphi) = \Delta_0 \left[\frac{2}{3} \sqrt{\frac{a}{3}} \cos(2\varphi) / (1 + a \cos^2(2\varphi))^{\frac{3}{2}} \right], \quad a > 1/2 \quad (12)$$

MUSRFIT theory line: `userFcn libGapIntegrals TGapNonMonDWave2 1 2 3 [4]`

(Parameters: T_c (K), Δ_0 (meV), a (1), a (1), [a_G (1)]. If a_G is given, the temperature dependence according to Eq.(7) will be used, otherwise Eq.(8) will be utilized.)

anisotropic s-wave gap [8]:

$$\Delta(\varphi) = \Delta_0 [1 + a \cos(4\varphi)], \quad 0 \leq a \leq 1 \quad (13)$$

MUSRFIT theory line: `userFcn libGapIntegrals TGapAnSWave 1 2 3 [4]`

(Parameters: T_c (K), Δ_0 (meV), a (1), [a_G (1)]. If a_G is given, the temperature dependence according to Eq.(7) will be used, otherwise Eq.(8) will be utilized.)

It is also possible to calculate a power law temperature dependence (in the two fluid approximation $n = 4$) and the dirty s-wave expression. Obviously for this no integration is needed.

Power law return function:

$$\frac{\lambda(0)^2}{\lambda(T)^2} = 1 - \left(\frac{T}{T_c}\right)^n \quad (14)$$

MUSRFIT theory line: `userFcn libGapIntegrals TGapPowerLaw 1 2`
 (Parameters: T_c (K), n (1))

dirty s-wave [9]:

$$\frac{\lambda(0)^2}{\lambda(T)^2} = \frac{\Delta(T)}{\Delta_0} \tanh \left[\frac{\Delta(T)}{2k_B T} \right] \quad (15)$$

with $\Delta(T)$ given by Eq.(??).

MUSRFIT theory line: `userFcn libGapIntegrals TGapDirtySWave 1 2 [3]`
 (Parameters: T_c (K), Δ_0 (meV), [a_G (1)]. If a_G is given, the temperature dependence according to Eq.(7) will be used, otherwise Eq.(8) will be utilized.)

Currently there are two gap functions to be found in the code which are *not* documented here: `TGapCosSqDWave` and `TGapSinSqDWave`. For details for these gap functions (superfluid density along the a/b -axis within the semi-classical model assuming a cylindrical Fermi surface and a mixed $d_{x^2-y^2} + s$ symmetry of the superconducting order parameter (effectively: $d_{x^2-y^2}$ with shifted nodes and a - b -anisotropy)) see the source code.

License

The GAPINTEGRALS library is free software; you can redistribute it and/or modify it under the terms of the GNU General Public License as published by the Free Software Foundation [10]; either version 2 of the License, or (at your option) any later version.

References

- [1] A. Suter, MUSRFIT *User Manual*, <https://wiki.intranet.psi.ch/MUSR/MusrFit>
- [2] T. Hahn, *Cuba - a library for multidimensional numerical integration*, Comput. Phys. Commun. **168** (2005) 78-95, <http://www.feynarts.de/cuba/>
- [3] R. Prozorov and R.W. Giannetta, *Magnetic penetration depth in unconventional superconductors*, Supercond. Sci. Technol. **19** (2006) R41-R67, and Erratum in Supercond. Sci. Technol. **21** (2008) 082003.
- [4] A. Carrington and F. Manzano, Physica C **385** (2003) 205
- [5] G. Deutscher, *Andreev-Saint-James reflections: A probe of cuprate superconductors*, Rev. Mod. Phys. **77** (2005) 109-135
- [6] H. Matsui *et al.*, *Direct Observation of a Nonmonotonic $d_{x^2-y^2}$ -Wave Superconducting Gap in the Electron-Doped High- T_c Superconductor $Pr_{0.89}LaCe_{0.11}CuO_4$* , Phys. Rev. Lett. **95** (2005) 017003
- [7] I. Eremin, E. Tsoncheva, and A.V. Chubukov, *Signature of the nonmonotonic d-wave gap in electron-doped cuprates*, Phys. Rev. B **77** (2008) 024508
- [8] ??
- [9] M. Tinkham, *Introduction to Superconductivity* 2nd ed. (Dover Publications, New York, 2004).
- [10] <http://www.gnu.org/licenses/old-licenses/gpl-2.0.html>